



Health Sciences and Human Services
Surgical Technology
HEALTH IMMUNIZATION RECORD

I _____ MD/NP/PAC
Type or print provider name *Provider signature*

Provider address *Provider telephone*

have examined _____ on _____
Name of student *Date of Exam*

For admission to **CSI SURGICAL TECHNOLOGY PROGRAM** and report the following immunization information:

The recommendations listed below follow the CDC recommendations for vaccinations for healthcare workers. **Healthcare workers include physicians, nurses, emergency medical personnel, dental professionals, and students, medical and nursing students, laboratory technicians, pharmacists, hospital volunteers, and administrative staff. (www.cdc.gov)**

Hepatitis B *REQUIRED* If you don't have documented evidence of a complete hepB vaccine series, or if you don't have an up-to-date blood test that shows you are immune to hepatitis B (i.e., no serologic evidence of immunity or prior vaccination) then you should: Get the 3-dose series (dose #1 now, #2 in 1 month, #3 approximately 5 months after #2). Get anti-HBs serologic tested 1–2 months after dose #3.

Tetanus and Diphtheria & Pertussis *REQUIRED* Get a one-time dose of Tdap as soon as possible if you have not received Tdap previously (regardless of when previous dose of Td was received). Get Td boosters every 10 years thereafter. Pregnant HCWs need to get a dose of Tdap during each pregnancy.

Measles, Mumps, Rubella *REQUIRED* If you were born in 1957 or later and have not had the MMR vaccine, or if you don't have an up-to-date blood test that shows you are immune to measles or mumps (i.e., no serologic evidence of immunity or prior vaccination), get 2 doses of MMR (1 dose now and the 2nd dose at least 28 days later).
If you were born in 1957 or later and have not had the MMR vaccine, or if you don't have an up-to-date blood test that shows you are immune to rubella, only 1 dose of MMR is recommended. However, you may end up receiving 2 doses, because the rubella component is in the combination vaccine with measles and mumps.

Varicella *REQUIRED* If you have not had chickenpox (varicella), if you haven't had varicella vaccine, or if you don't have an up-to-date blood test that shows you are immune to varicella (i.e., no serologic evidence of immunity or prior vaccination) get 2 doses of varicella vaccine, 4 weeks apart.

Meningococcal *REQUIRED* Those who are routinely exposed to isolates of *N. meningitidis* should get one dose.

Influenza (Flu) *REQUIRED* Get 1 dose of influenza vaccine annually during flu season.

Please complete immunization record on following pages.

